



RAINFAST
WITHIN
ONE HOUR

ERASER MAX


SUPER CONCENTRATE

Controls Grasses, Broadleaf Weeds, Brush
and Vines for up to One Year.

Not for use near desirable vegetation. For outdoor use only.



KUDZU



DALLISGRASS



THISTLE

ACTIVE INGREDIENTS:

Glyphosate, isopropylamine salt * **43.68%**

Imazapyr, isopropylamine salt * * **00.78%**

OTHER INGREDIENTS: **55.54%**

TOTAL: **100.00%**

ACID EQUIVALENT: * Glyphosate: 3.158 lbs/gal

** Imazapyr: 0.062 lbs/gal

KEEP OUT OF REACH OF CHILDREN DANGER - PELIGRO

Si usted no entiende la etiqueta, busque a alguien para que se
la explique a usted en detalle. (If you do not understand the label,
find someone to explain it to you in detail.)

See booklet for First Aid and additional Precautionary Statements



0 72693 24885 4

EPA Reg. No. 35935-94-53883

EPA Est. No. 88746-GA-01

NET CONTENTS:

ONE QUART (946 mL)

PRECAUTIONARY STATEMENT
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
DANGER - PELIGRO

Corrosive. Causes irreversible eye damage. Harmful if swallowed. Do not get in eyes or on clothing.

PERSONAL PROTECTIVE EQUIPMENT (PPE):

Some materials that are chemical-resistant to this product are natural rubber > 14 mils. If you want more options, follow the instructions for category A on an EPA chemical-resistance selection chart.

Mixers, loaders, applicators, and other handlers must wear:

- Long-sleeve shirt and long pants,
- Shoes plus socks,
- Chemical resistant gloves for all mixers and loaders, plus applicators using handheld equipment,
- Protective eyewear (goggles, face shield, or safety glasses).

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry. Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands with plenty of soap and water before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product.
- Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

FIRST AID

IF IN EYES

- Hold eye open and rinse slowly and gently with water for 15 to 20 minutes.
- Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
- Call a poison control center or doctor for treatment advice.

IF SWALLOWED

- Call a poison control center or doctor immediately for treatment advice.
- Have person sip a glass of water if able to swallow.
- Do not induce vomiting unless told to by a poison control center or doctor.
- Do not give anything to an unconscious person.

HOT LINE NUMBER

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For information on this product, contact the National Pesticide Information Center, 1-800-858-7378, Monday-Friday, 8:00 AM-12:00 PM PST, or the SafetyCall HotLine at 1-866-897-8050. You may also contact the National Poison Control Center, 1-800-222-1222, day or night, for emergency medical treatment information. **FOR CHEMICAL EMERGENCY:** Spill, leak, fire, exposure, or accident call CHEMTREC 1-800-424-9300.

ENVIRONMENTAL HAZARDS

This product is toxic to plants. Drift and run-off may be hazardous to plants in water adjacent to treated areas. Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater or rinsate. See Directions for Use for additional precautions and requirements.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Read the entire label before using this product. Use strictly in accordance with label precautionary statements and directions.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

Do not enter or allow others to enter treated areas until sprays have dried.

Do not use on feed or food crops.

Keep people and pets out of the area during application.

Applicators are required to use a nozzle height below 4 feet above the ground or plant canopy and coarse or coarser droplet size (ASABE S572) or, if specifically using a spinning atomizer nozzle, applicators are required to use a volume mean diameter (VMD) of 385 microns or greater. Applications with wind speeds greater than 10 mph are prohibited. Applications into temperature inversions are prohibited.

WHERE TO USE

Use on non-crops areas: Industrial sites, Petroleum tank farms, Lumberyards, Manufacturing Sites, Storage Areas, Airports, Roadsides, Curbs, Sidewalks, Driveways, Parking areas, Brick walks, Gravel paths, Patios, Vacant lots, Paved areas (including crack and crevice), Warehouse areas, Urban areas, Recreational areas, Play areas, Tennis courts, Fencerows, Along fences of paddocks, corrals, feedlots, and livestock holding areas, Hedgerows, Dry Ditchbanks, Around domestic dwellings, Around farm buildings and barns, Agricultural uncultivated areas, Shelter beds, Farm yards, Farm premises, Outdoor buildings, structures and shelters such as beef barns, hog barns, dairy barns, stables, dog kennels and poultry houses, and For establishment and maintenance of wildlife openings

Apply to locations only where no vegetation growth is desired for up to one year.

WHEN TO USE

Apply to bare ground or when weeds are present. If weeds are taller than six inches, mow or cut down weeds and remove debris before treatment, and allow for 2 weeks of new growth. If woody vines or brush is mowed or cut, wait until full leaf formation before treatment.

To prevent growth of weeds under asphalt driveways, play areas, tennis courts and parking areas, apply this product before laying surfacing materials.

Treat area uniformly to maximize results.

Re-treatment every 6 to 8 months may be necessary in areas of heavy rainfall.

RESTRICTIONS

- Apply on a sunny day with temperatures above 60°F.
- Do not apply to or allow product to drift onto desirable plants.
- Only apply to areas where no vegetation is desired for one year.
- Avoiding product runoff to nearby desirable plants.
- Do not apply near bodies of water including, fish pools, ponds, lakes or streams.
- Do not apply over the roots systems or desirable plants. For trees and shrubs do not apply closer than twice the distance from the trunk to the drip line as roots may be within this area.
- Do not apply to slopes in landscape as movement on soil surfaces may damage desirable plants down slope.
- Do not apply next to a fence if desirable vegetation is growing on the other side, or if future planting is intended.
- Do not use before planting lawns, fruits, vegetables, flowers, or other plants because this product remains active in soil for up to one year.
- In soils with heavy clay, weed control may exceed one year.
- Do not use on desirable lawns.
- Do not apply when wind speeds are greater than 10 miles per hour.
- Do not apply to powdery, dry or light sandy soils when the soil may be blown by wind. If necessary soil surface should be settled by rain or irrigation.

HOW TO APPLY

Total Vegetation Control (Bareground)

Use hand operated spray equipment such as backpack sprayers, pump-up sprayers or sprinkling cans, or ATV/Tractor mounted sprayers. Mix 7.4 fl. oz. (0.46 pints) of product with water to produce 1 to 10 gallons of spray solution. Optimum performance will be achieved from applications to actively growing weeds in early summer.

Uniformly spray the mix over areas where total vegetation control and bareground is desired. 7.4 fl. oz. of product within 1 to 10 gallons of spray solution should be applied to 1,000 square feet.

Maximum application rate of this product is 2.5 gallons per acre per year (7.4 fl. oz. per 1,000 sq. ft. per year).

Spot Control of Brush, Vines, and Weeds

Use hand operated spray equipment such as backpack sprayers, pump-up sprayers or sprinkling cans, or ATV/Tractor mounted sprayers. Mix 6 oz. of product per 1 gallon of spray solution. Thoroughly and uniformly spray foliage of woody plants to wet. For optimum performance:

- Apply to actively growing brush, vines and weeds.
- Woody brush and vines are controlled better in late summer and fall after fruit formation.
- If woody vines and brush are mowed or cut, wait until full leaf formation before treatment.
- Do not cut or mow brush, vines or perennial weeds for 2 months following treatment.

Maximum application rate of this product is 2.5 gallons per acre per year (7.4 fl. oz. per 1,000 sq. ft. per year).

Weeds Controlled lists

GRASS/NUTSEGE WEED SPECIES CONTROLLED

Bahiagrass

Paspalum notatum

Bentgrass

Agrostis spp.

Bermudagrass

Cynodon dactylon

Bluegrass, Kentucky

Poa spp.

Brome

Bromus spp.

Bromegrass, smooth

Bromus inermis

Canarygrass, reed

Phalaris arundinacea

Cattail

Typha spp.

Cogongrass

Imperata cylindrical

Dallisgrass

Paspalum dilatatum

Fescues

Festuca spp.

Fescue, tall

Festuca arundinacea

Guineagrass

Panicum maximum

Johnsongrass

Sorghum halepense

Kikuyugrass

Pennisetum clandestinum

Knotgrass

Paspalum distichum

Muhly, wirestem

Muhlenbergia frondosa

Napier grass

Pennisetum purpureum

Nutsedge; purple, yellow

Cyperus rotundus

Cyperus esculentus

Orchardgrass

Dactylis glomerata

Pampas grass

Cortaderia spp.

Panicum

Panicum spp.

GRASS/NUTSEGE WEED SPECIES CONTROLLED

(continued)

Para grass

Brachiaria mutica

Phragmites*

Phragmites spp.

Quackgrass

Agropyron repens

Reed, giant

Arundo donax

Ryegrass, perennial

Lolium perenne

Timothy

Phleum pratense

Torpedograss*

Panicum repens

Vaseygrass

Paspalum urvillei

Velvetgrass

Holcus spp.

Wheatgrass, western

Agropyron smithii

* Partial Control

BROADLEAF WEED SPECIES CONTROLLED

Alfalfa

Medicago sativa

Alligatorweed*

Alternanthera philoxeroides

Anise (fennel)

Foeniculum vulgare

Artichoke, Jerusalem

Helianthus tuberosus

Bassia, fivehook

Bassia hyssopifolia

Bindweed, field

Convolvulus arvensis

Blueweed, Texas

Helianthus ciliaris

Bracken fern

Pteridium aquilinum

Bursage, woollyleaf

Franseria tomentosa

Clover, red

Trifolium pretense

Clover, white

Trifolium repens

Dandelion

Taraxacum officinale

Dock, curly

Rumex crispus

Dogbane, hemp

Apocynum cannabinum

Fiddleneck

Amsinckia spp.

Fleabane

Erigeron spp.

Fleabane, hairy

Conyza bonariensis

Horsenettle

Solanum carolinense

Horseradish

Armoracia rusticana

Ice plant

Mesembryanthemum crystallinum

Knapweed

Centaurea repens

Kochia

Kochia scoparia

Lantana

Lantana camara

Lespedeza

Lespedeza spp.

Lettuce, prickly

Lactuca serriola

Milkweed

Asclepias spp.

Mullein, common

Verbascum thapsus

* Partial Control

BROADLEAF WEED SPECIES CONTROLLED

(continued)

Nightshade, silverleaf

Solanum elaeagnifolium

Poison hemlock

Conium maculatum

Ragweed, common

Ambrosia artemisiifolia

Ragweed, giant

Ambrosia trifida

Smartweed, Pennsylvania

Polygonum pennsylvanicum

Smartweed, swamp

Polygonum coccineum

Sowthistle, annual

Sonchus oleraceus

Spurge, leafy*

Euphorbia esula

Starthistle, yellow

Centaurea solstitialis

Sunflower

Helianthus annuus

Thistle, artichoke

Cynara cardunculus

Thistle, Canada

Cirsium arvense

Thistle, Russian

Salsola kali

Velvetleaf

Abutilon theophrasti

* Partial Control

TREE, BRUSH AND VINE SPECIES CONTROLLED

Alder

Alnus spp.

Ash

Fraxinum spp.

Aspen, quaking

Populus tremuloides

Balsam apple*

Momordica charantia

Bearmat (Bearclover)

Chamaebatia foliolosa

Beech

Fagus grandifolia

Birch

Betula spp.

Blackberry

Rubus spp.

Blackgum

Nyssa spp.

Bracken

Peridium spp.

Broom:

French

Cytisus monspessulanus

Scotch

Cytisus scoparius

Buckwheat, California*

Eriogonum fasciculatum

Cascara*

Rhamnus purshiana

Catsclaw*

Acacia greggi

Ceanothus*

Ceanothus spp.

Chamise

Adenostoma fasciculatum

Cherry:

Bitter

Prunus emarginata

Black

Prunus serotina

Pin

Prunus pennsylvanica

Coyote brush

Baccharis consanguinea

Creeper, Virginia*

Parthenocissus quinquefolia

Dewberry

Rubus trivialis

*Partial Control, **This product is not registered in California for use on Russian olive.

TREE, BRUSH AND VINE SPECIES CONTROLLED

(continued)

Dogwood*

Cornus spp.

Elderberry

Sambucus spp.

Elm*

Ulmus spp.

Eucalyptus

Eucalyptus spp.

Gorse

Ulex europaeus

Hasardia*

Haplopappus squamosus

Hawthorn

Crataegus spp.

Hazel

Corylus spp.

Hickory*

Carya spp.

Holly, Florida/Brazilian

Peppertree*

Schinus terebinthifolius

Honeysuckle

Lonicera spp.

Hornbeam, American*

Carpinus caroliniana

Kudzu

Pueraria lobata

Locust, black*

Robinia pseudoacacia

Madrone

Arbutus menziesii

Manzanita

Arctostaphylos spp.

Maple:

Red**

Acer rubrum

Sugar

Acer saccharum

Vine

Acer circinatum

Monkey Flower*

Mimulus guttatus

Oak:

Black*

Quercus velutina

Northern Pin

Quercus palustris

Post

Quercus stellata

Red

Quercus rubra

Southern Red

Quercus falcata

White*

Quercus alba

Persimmon*

Diospyros spp.

Pine

Pinus spp.

Poison Ivy

Rhus radicans

Poison Oak

Rhus toxicodendron

Poplar, yellow*

Liriodendron tulipifera

Raspberry

Rubus spp.

Redbud, eastern

Cercis canadensis

Redvine*

Brunnichia ovata

Rose, multiflora

Rosa multiflora

Russian olive**

Elaeagnus angustifolia

Sage: black, white

Salvia spp.

Sagebrush, California

Artemisia californica

Salmonberry

Rubus spectabilis

*Partial Control, **This product is not registered in California for use on Russian olive.

TREE, BRUSH AND VINE SPECIES CONTROLLED

(continued)

Salt cedar

Tamarix spp.

Sassafras

Sassafras aibidum

Sourwood

Oxydendrum arboreum

Sweet potato, wild*

Ipomoea pandurata

Sumac:

Poison*

Rhus vernix

Smooth*

Rhus glabra

Winged*

Rhus copallina

Sweetgum

Liquidambar styraciflua

Swordfern*

Polystichum munitum

Tallowtree, Chinese

Sapium sebiferum

Tan Oak

Lithocarpus densiflorus

Thimbleberry

Rubus parviflorus

Tobacco, tree*

Nicotiana glauca

Trumpetcreeper*

Campsis radicans

Waxmyrtle, southern*

Myrica cerifera

Willow

Salix spp.

*Partial Control, **This product is not registered in California for use on Russian olive.

STORAGE AND DISPOSAL

Do not contaminate water, foodstuffs, feed or seed by storage or disposal.

PESTICIDE STORAGE: Store product in original container in a safe place and protect from freezing.

PESTICIDE DISPOSAL: Pesticide wastes may be acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

STORAGE AND DISPOSAL (continued)

CONTAINER DISPOSAL:

Nonrefillable containers smaller than 5 gallons:

Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available. Triple rinse container (or equivalent) promptly after emptying. **Triple rinse as follows:** Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by State and local authorities. Plastic containers are also disposable by incineration, or, if allowed by State and local authorities, by burning. If burned stay out of smoke.

WARRANTY STATEMENT

Control Solutions, Inc. warrants that this product conforms to the chemical description on the label thereof and is reasonably fit for purposes stated on such label only when used in accordance with directions under normal use conditions. It is impossible to eliminate all risks inherently associated with use of this product. Crop injury, ineffectiveness, or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Control Solutions, Inc. To the extent allowed by applicable law, Control Solutions, Inc. shall not be liable for consequential, special, or indirect damages resulting from the use or handling of this product. All such risks shall be assumed by the Buyer. In addition to the foregoing, no purchaser of this product (other than an end user) shall be entitled to any reimbursement for any loss suffered as a result of any suspension or cancellation of the registration for this product by the U.S. Environmental Protection Agency. Except, as expressly provided herein and to the extent consistent with applicable law, Control Solutions, Inc. makes no warranties, guarantees, or representations of any kind, either expressed or implied, or by usage of trade, statutory or otherwise, with regard to the product sold, including, but not limited to merchantability, fitness for a particular purpose, use or eligibility of the product for any

particular trade usage. The exclusive remedy of any buyer or user of this product for any and all losses, injuries, or damage resulting from or in any way arising from the use, handling, or application of this product, whether in contract, warranty, tort, negligence, strict liability, or otherwise, shall be damages not exceeding the purchase price paid for this product or, at Control Solutions, Inc. election, the replacement of this product.

Buyer and user assume all responsibilities for safety and use not in strict accordance with directions.

RV121211

Manufactured for
CONTROL SOLUTIONS, INC.
5903 Genoa-Red Bluff
Pasadena, TX 77507-1041



ERASER MAX SUPER CONCENTRATE

Not for use near desirable vegetation. For outdoor use only.

- **FAST ACTING- one application kills weeds for up to 1 year**
- **Kills weeds and grasses, roots and all**
- **Use 7.4 fl. oz. per 1,000 sq. ft. per year**
- **Season long control**
- **Rainfast within one hour**

ACTIVE INGREDIENTS:

Glyphosate, isopropylamine salt*	43.68%
Imazapyr, isopropylamine salt**	00.78%
OTHER INGREDIENTS:	55.54%
TOTAL:	100.00%

ACID EQUIVALENT: * Glyphosate: 3.158 lbs/gal ** Imazapyr: 0.062 lbs/gal

EPA Reg. No. 35935-94-53883

EPA Est. No. 88746-GA-01

KEEP OUT OF REACH OF CHILDREN DANGER – PELIGRO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
(If you do not understand the label, find someone to explain it to you in detail.)

See booklet for First Aid and additional Precautionary Statements

FIRST AID

IF IN EYES	<ul style="list-style-type: none">• Hold eye open and rinse slowly and gently with water for 15 to 20 minutes.• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.• Call a poison control center or doctor for treatment advice.
IF SWALLOWED	<ul style="list-style-type: none">• Call a poison control center or doctor immediately for treatment advice.• Have person sip a glass of water if able to swallow.• Do not induce vomiting unless told to by a poison control center or doctor.• Do not give anything to an unconscious person.

HOT LINE NUMBER

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For information on this product, contact the National Pesticide Information Center, 1-800-858-7378, Monday-Friday, 8:00 AM-12:00 PM PST, or the SafetyCall HotLine at 1-866-897-8050. You may also contact the National Poison Control Center, 1-800-222-1222, day or night, for emergency medical treatment information.

FOR CHEMICAL EMERGENCY: Spill, leak, fire, exposure, or accident call CHEMTREC 1-800-424-9300.
For Medical Emergencies Only, Call (866) 897-8050



CONTROL SOLUTIONS INC.

5903 GENOA-RED BLUFF ROAD
PASADENA TEXAS 77507

Net Contents: 1 QUART (946 mL)