# BONANZA™ Fly Bait

- For the Control of House Flies In and Around Areas Such as Livestock Facilities and Stables, Warehouses, Dumpsters, Recycling Areas, and Outside Non-Food Contact Areas of Restaurants, Bakeries, Supermarkets, Grease Pits, or Loading Docks.
- For use in scatter, bait station and paint on applications

# Callout circles

- Treats 1,333 square feet [For ½# package size]
- Treats 10,668 square feet [for 4# package size]
- Treats 101.346 square feet [for 38# package size]

# **ACTIVE INGREDIENT:**

Imidacloprid; 1-[6-Chloro-3-pyridinyl)methyl]-N nitro-2-imidazolidinimine	0.50%
Z-9-Tricosene	0.10%
OTHER INGREDIENTS	99.4%
TOTAL	100.00%

# STOP-Read the label before use. KEEP OUT OF REACH OF CHILDREN CAUTION

FIRST AID	
ment advice.	
control center or doctor.	
nutes.	
give artificial respiration,	
advice.	
auvice.	
5-20 minutes.	
s, then continue rinsing eye.	
5-20 minutes.	

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact SafetyCall® International (866) 897-8050 for emergency medical treatment information. Note to Physician: No specific antidote is available. Treat the patient symptomatically.

# PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

**CAUTION:** Harmful if swallowed, absorbed through skin, or inhaled. Avoid contact with eyes, skin or clothing. Wash thoroughly with soap and water after handling and before eating, drinking or using tobacco.

# **ENVIRONMENTAL HAZARDS**

This product is highly toxic to aquatic invertebrates. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters or rinsate. This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area. This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

# STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

Pesticide Storage: Store in original container in a dry place. Store in locked area out of reach of children.

Pesticide Disposal: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spilled bait, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for Guidance.

**Container Disposal:** Non-refillable container. Do not reuse or refill this container. Empty container through use according to label instructions. Offer for recycling, if available. If recycling is not available; then dispose of container in a sanitary landfill or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

**BONANZA FLY BAIT** is a fast acting ready-to-use house fly bait that may be applied as scatter bait, paint-on, or in bait stations.

It contains a specially formulated mixture of attractants in extruded pellet form and it works against house flies which are resistant to organophosphates and carbamates.

**BONANZA FLY BAIT** is a ready-to-use bait containing 0.5% imidacloprid and the attractant Z-9-Tricosene. It also contains Bitrex® bittering agent to help prevent ingestion by animals, children and pets.

BONANZA FLY BAIT controls nuisance house flies

#### COVERAGE

1.5 oz./250 sq ft (1.5 oz = 3 Tablespoons)

#### RE-APPLY

Re-apply every 7 days as needed.

### **DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. IMPORTANT: Read entire label and Conditions of Sale before using this product.

# HOUSE FLY MANAGEMENT

WHERE TO APPLY

Do not place BONANZA FLY BAIT in areas accessible to animals, children and pets.

# **Scatter Application**

For use around the **outside** of:

- Agricultural production facilities like broiler houses, turkey houses, caged layer houses, feedlots, dairy barns, and swine confinement buildings.
- Canneries; dairy, meat and poultry processing plants; food processing plants, warehouses; dumpsters; recycling areas and loading docks.

# **Scatter Application**

For use in walkways of:

• Caged layer houses.

# **Bait Station Application**

For use **inside** of:

- Horse Stables
- Broiler, turkey, and caged layer houses.
- Swine confinement buildings.
- Dairy barns and milking parlors.

## Paint on Application

- For use inside and around the outside of.
- Broiler, turkey, and caged layer houses.
- Swine confinement buildings.
- Dairy barns.

# **USE RATE**

• 1.5 oz/250 sq ft (scatter bait and bait station).

### **HOW TO APPLY**

Apply the bait in many locations. Avoid cool or windy locations because house flies prefer warm, motionless sites for resting. Time applications to begin at the start of the season before house fly populations have reached their peak.

- Scatter Application: Scatter the bait directly from the container onto dry level surfaces so that the individual granules lie near each other without forming small piles.
- Bait Station Application: Place in any fly bait station. Use 1 bait station to cover approximately 250 sq. ft., and add up to 1.5 oz of bait per station. Secure bait stations at least 4 feet above the ground.
- Paint-on Application: Mix 1.5 oz. bait with 1 fl oz warm water (3 lbs in 1 quart water) and stir thoroughly. Let stand for about 15 minutes until a paste consistency suitable for painting has formed. Apply the paste with a brush to surfaces where house flies rest, avoiding surfaces that are dusty.

### RESTRICTIONS

- Do not place bait in areas accessible to food producing animals, children and pets.
- Do not allow contact with food or water.
- Use environmental controls, such as proper light management.
- Do not apply directly onto milking equipment.
- Do not apply inside food processing plants or restaurants.

# **MANAGEMENT TIPS**

#### For Best Results:

- Use the bait as part of an overall Integrated Pest Management program.
- Use proper sanitation and manure management.
- Use along with dilutable sprays.
- Use environmental controls, such as proper light management.
- Consult with Cooperative Extension Services.

NOTE: CONTROL SOLUTIONS, INC. WARRANTS THAT THIS PRODUCT CONFORMS TO THE CHEMICAL DESCRIPTION ON THE LABEL AND IS REASONABLY FIT FOR THE PURPOSES STATED ON THE LABEL WHEN USED IN ACCORDANCE WITH THE DIRECTIONS FOR USE. IT IS IMPOSSIBLE TO ELIMINATE ALL RISKS INHERENTLY ASSOCIATED WITH THE USE OF THIS PRODUCT. INEFFECTIVENESS OR OTHER UNINTENDED CONSEQUENCES MAY RESULT BECAUSE OF SUCH FACTORS AS WEATHER CONDITIONS, PRESENCE OF OTHER MATERIALS, OR THE MANNER OR APPLICATION, ALL OF WHICH ARE BEYOND THE CONTROL OF CONTROL SOLUTIONS, INC. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, CONTROL SOLUTIONS, INC. SHALL NOT BE LIABLE FOR CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. ALL SUCH RISKS SHALL BE ASSUMED BY THE BUYER

EPA REG. NO. 53883-238 EPA EST. NO. 44616-MO-1(A) 53883-TX-002 (B)

Circled letter is first letter of lot number.

Control Solutions, Inc. 5903 Genoa-Red Bluff Pasadena, TX 77507-1041